

Fitting genotype by environment models in sommer

Giovanny Covarrubias-Pazaran

2020-04-09

The sommer package was developed to provide R users a powerful and reliable multivariate mixed model solver. The package is focused in problems of the type $p > n$ (more effects to estimate than observations) and its core algorithm is coded in C++ using the Armadillo library. This package allows the user to fit mixed models with the advantage of specifying the variance-covariance structure for the random effects, and specify heterogeneous variances, and obtain other parameters such as BLUPs, BLUEs, residuals, fitted values, variances for fixed and random effects, etc.

The purpose of this vignette is to show how to fit different genotype by environment (GxE) models using the sommer package:

- 1) Single environment model
- 2) Multienvironment model: Main effect model
- 3) Multienvironment model: Diagonal model (DG)
- 4) Multienvironment model: Compound symmetry model (CS)
- 5) Multienvironment model: Compound symmetry + diagonal model (CS+DG)
- 6) Multienvironment model: Unstructured model (US)
- 7) Multienvironment model: Random regression model (RR)
- 8) Multienvironment model: Other covariance structures for GxE

When the breeder decides to run a trial and apply selection in a single environment whether because the amount of seed is a limitation or there's no availability for any location the breeder takes the risk of selecting material for a target population of environments (TPEs) and this environment tested not being representative of the larger TPE. Therefore, many breeding programs try to base their selection decision using multi-environment trial (MET) data. Although, models could be adjusted by adding additional information like spatial information, experimental design information, etc., in this tutorial we will focus mainly on the covariance structures for GxE and the incorporation of relationship matrices for the genotype effect.

1) Single environment model

A single environment model is the one that is fitted when the breeding program can only afford one location leaving out the possible information available from other environments. This will be used to further expand to GxE models.

```
library(sommer)
data(DT_example)
DT <- DT_example
A <- A_example

ansSingle <- mmer(Yield~1,
                     random= ~ vs(Name, Gu=A),
                     rcov= ~ units,
                     data=DT)

## iteration      LogLik      wall      cpu(sec)      restrained
```

```

##      1    -80.9858  21:52:12      0      0
##      2    -79.2137  21:52:12      0      0
##      3    -78.8346  21:52:12      0      0
##      4    -78.8088  21:52:12      0      0
##      5    -78.8087  21:52:12      0      0

summary(ansSingle)

## =====
## Multivariate Linear Mixed Model fit by REML
## **** sommer 4.1 ****
## =====
##      logLik      AIC      BIC Method Converge
## Value -78.80875 159.6175 162.8378      NR      TRUE
## =====
## Variance-Covariance components:
##          VarComp VarCompSE Zratio Constraint
## u:Name.Yield-Yield   6.529     2.202  2.965  Positive
## units.Yield-Yield   13.868     1.633  8.494  Positive
## =====
## Fixed effects:
##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept)  11.74     0.4876   24.07
## =====
## Groups and observations:
##   Yield
## u:Name   41
## =====
## Use the '$' sign to access results and parameters

```

In this model the only term to be estimated is the one for the germplasm (here called Name). For the sake of example we have added a relationship matrices among the levels of the random effect Name. This is just a diagonal matrix with as many rows and columns as levels present in the random effect Name, but any other non-diagonal relationship matrix could be used.

2) MET: main effect model

A multi environment model is the one that is fitted when the breeding program can afford more than one location. This assumes that GxE doesn't exist and that the main genotype effect plus the fixed effect for environment is enough to predict the genotype effect in all locations of interest.

```

ansMain <- mmer(Yield~Env,
                  random= ~ vs(Name, Gu=A),
                  rcov= ~ units,
                  data=DT)

## iteration      LogLik      wall    cpu(sec)  restrained
##      1    -36.8096  21:52:12      0          0
##      2    -33.211   21:52:12      0          0
##      3    -32.6234  21:52:12      0          0
##      4    -32.5942  21:52:12      0          0
##      5    -32.5942  21:52:12      0          0

summary(ansMain)

## =====

```

```

## Multivariate Linear Mixed Model fit by REML
## **** sommer 4.1 ****
## =====
##      logLik      AIC      BIC Method Converge
## Value -32.59421 71.18842 80.84949     NR     TRUE
## =====
## Variance-Covariance components:
##           VarComp VarCompSE Zratio Constraint
## u:Name.Yield-Yield   4.856    1.5233  3.188  Positive
## units.Yield-Yield    8.109    0.9615  8.434  Positive
## =====
## Fixed effects:
##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept) 16.385    0.5849 28.012
## 2 Yield EnvCA.2012  -5.688    0.5741 -9.908
## 3 Yield EnvCA.2013  -6.218    0.6107 -10.182
## =====
## Groups and observations:
##   Yield
## u:Name    41
## =====
## Use the '$' sign to access results and parameters

```

3) MET: diagonal model (DG)

A multi environment model is the one that is fitted when the breeding program can afford more than one location. This assumes that GxE doesn't exist and that the main genotype effect plus the fixed effect for environment is enough to predict the genotype effect in all locations of interest.

```

ansDG <- mmer(Yield~Env,
                 random= ~ vs(ds(Env),Name, Gu=A),
                 rcov= ~ units,
                 data=DT)

## iteration  LogLik    wall  cpu(sec)  restrained
##   1       -42.26  21:52:12      0          0
##   2       -26.3735 21:52:12      0          0
##   3       -21.5756 21:52:12      0          0
##   4       -21.05   21:52:12      0          0
##   5       -21.0417 21:52:12      0          0
##   6       -21.0416 21:52:13      1          0

summary(ansDG)

## =====
## Multivariate Linear Mixed Model fit by REML
## **** sommer 4.1 ****
## =====
##      logLik      AIC      BIC Method Converge
## Value -21.04157 48.08315 57.74421     NR     TRUE
## =====
## Variance-Covariance components:
##           VarComp VarCompSE Zratio Constraint
## CA.2011:Name.Yield-Yield   17.493    6.1099  2.863  Positive
## CA.2012:Name.Yield-Yield    5.337    1.7662  3.022  Positive

```

```

## CA.2013:Name.Yield-Yield    7.884    2.5526  3.089   Positive
## units.Yield-Yield          4.381    0.6493  6.747   Positive
## =====
## Fixed effects:
##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept) 16.621     0.948  17.532
## 2 Yield EnvCA.2012   -5.958     1.045  -5.699
## 3 Yield EnvCA.2013   -6.662     1.098  -6.067
## =====
## Groups and observations:
##   Yield
## CA.2011:Name    41
## CA.2012:Name    41
## CA.2013:Name    41
## =====
## Use the '$' sign to access results and parameters

```

4) MET: compound symmetry model (CS)

A multi environment model is the one that is fitted when the breeding program can afford more than one location. This assumes that GxE doesn't exist and that the main genotype effect plus the fixed effect for environment is enough to predict the genotype effect in all locations of interest.

```

E <- diag(length(unique(DT$Env)))
rownames(E) <- colnames(E) <- unique(DT$Env)
EA <- kronecker(E,A, make.dimnames = TRUE)
ansCS <- mmmer(Yield~Env,
                 random= ~ vs(Name, Gu=A) + vs(Env:Name, Gu=EA),
                 rcov= ~ units,
                 data=DT)

```

```

## iteration  LogLik    wall  cpu(sec)  restrained
##   1       -31.2668 21:52:13    0        0
##   2       -23.2804 21:52:13    0        0
##   3       -20.4746 21:52:13    0        0
##   4       -20.1501 21:52:13    0        0
##   5       -20.1454 21:52:13    0        0
##   6       -20.1454 21:52:13    0        0

```

```
summary(ansCS)
```

```

## =====
##           Multivariate Linear Mixed Model fit by REML
## ***** sommer 4.1 *****
## =====
##   logLik      AIC      BIC Method Converge
## Value -20.14538 46.29075 55.95182    NR    TRUE
## =====
## Variance-Covariance components:
##                               VarComp VarCompSE Zratio Constraint
## u:Name.Yield-Yield      3.682     1.691  2.177   Positive
## u:Env:Name.Yield-Yield  5.173     1.495  3.460   Positive
## units.Yield-Yield       4.366     0.647  6.748   Positive
## =====
## Fixed effects:

```

```

##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept)  16.496    0.6855  24.065
## 2 Yield EnvCA.2012  -5.777    0.7558  -7.643
## 3 Yield EnvCA.2013  -6.380    0.7960  -8.015
## -----
## Groups and observations:
##       Yield
## u:Name      41
## u:Env:Name  123
## -----
## Use the '$' sign to access results and parameters

```

5) MET: compound symmetry plus diagonal (CS+DIAG)

A multi environment model is the one that is fitted when the breeding program can afford more than one location. This assumes that GxE doesn't exist and that the main genotype effect plus the fixed effect for environment is enough to predict the genotype effect in all locations of interest.

```

ansMain <- mmmer(Yield~Env,
                    random= ~ vs(Name, Gu=A) + vs(ds(Env),Name, Gu=A),
                    rcov= ~ units,
                    data=DT)

```

```

## iteration  LogLik     wall    cpu(sec)  restrained
##   1      -31.2668  21:52:13      0          0
##   2      -21.0887  21:52:13      0          0
##   3      -18.4752  21:52:13      0          0
##   4      -18.1673  21:52:13      0          0
##   5      -18.1618  21:52:13      0          0
##   6      -18.1616  21:52:13      0          0

```

```
summary(ansMain)
```

```

## -----
##               Multivariate Linear Mixed Model fit by REML
## **** sommer 4.1 ****
## -----
##      logLik      AIC      BIC Method Converge
## Value -18.16164 42.32327 51.98434      NR      TRUE
## -----
## Variance-Covariance components:
##           VarComp VarCompSE Zratio Constraint
## u:Name.Yield-Yield      2.965    1.5055  1.969  Positive
## CA.2011:Name.Yield-Yield 10.424    4.4544  2.340  Positive
## CA.2012:Name.Yield-Yield  2.658    1.8032  1.474  Positive
## CA.2013:Name.Yield-Yield  5.702    2.5113  2.271  Positive
## units.Yield-Yield        4.398    0.6517  6.748  Positive
## -----
## Fixed effects:
##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept)  16.511    0.8269  19.967
## 2 Yield EnvCA.2012  -5.809    0.8593  -6.760
## 3 Yield EnvCA.2013  -6.423    0.9358  -6.864
## -----
## Groups and observations:

```

```

##          Yield
## u:Name      41
## CA.2011:Name 41
## CA.2012:Name 41
## CA.2013:Name 41
## =====
## Use the '$' sign to access results and parameters

```

6) MET: unstructured model (US)

A multi environment model is the one that is fitted when the breeding program can afford more than one location. This assumes that GxE doesn't exist and that the main genotype effect plus the fixed effect for environment is enough to predict the genotype effect in all locations of interest.

```

ansUS <- mmmer(Yield~Env,
                 random= ~ vs(us(Env),Name, Gu=A),
                 rcov= ~ units,
                 data=DT)

## iteration   LogLik    wall   cpu(sec)  restrained
##   1       -37.9059  21:52:13      0          0
##   2       -19.0506  21:52:13      0          0
##   3       -14.6786  21:52:13      0          0
##   4       -14.2203  21:52:13      0          0
##   5       -14.2098  21:52:13      0          0
##   6       -14.2095  21:52:13      0          0

summary(ansUS)

## =====
## Multivariate Linear Mixed Model fit by REML
## **** sommer 4.1 ****
## =====
##      logLik      AIC      BIC Method Converge
## Value -14.20951 34.41901 44.08008     NR      TRUE
## =====
## Variance-Covariance components:
##                                         VarComp VarCompSE Zratio Constraint
## CA.2011:Name.Yield-Yield        15.994   5.381  2.972  Positive
## CA.2012:CA.2011:Name.Yield-Yield  6.172   2.503  2.465 Unconstr
## CA.2012:Name.Yield-Yield       5.273   1.750  3.013  Positive
## CA.2013:CA.2011:Name.Yield-Yield  6.366   3.069  2.074 Unconstr
## CA.2013:CA.2012:Name.Yield-Yield  0.376   1.535  0.245 Unconstr
## CA.2013:Name.Yield-Yield        7.689   2.490  3.088  Positive
## units.Yield-Yield                4.386   0.650  6.748  Positive
## =====
## Fixed effects:
##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept) 16.341    0.8141  20.072
## 2 Yield EnvCA.2012  -5.696    0.7406  -7.692
## 3 Yield EnvCA.2013  -6.286    0.8202  -7.664
## =====
## Groups and observations:
##          Yield
## CA.2011:Name 41

```

```

## CA.2012:CA.2011:Name    82
## CA.2012:Name            41
## CA.2013:CA.2011:Name    82
## CA.2013:CA.2012:Name    82
## CA.2013:Name            41
## =====
## Use the '$' sign to access results and parameters
# adjust variance BLUPs by adding covariances
# ansUS$U[1:6] <- unsBLUP(ansUS$U[1:6])

```

7) MET: random regression model

A multi environment model is the one that is fitted when the breeding program can afford more than one location. This assumes that GxE doesn't exist and that the main genotype effect plus the fixed effect for environment is enough to predict the genotype effect in all locations of interest.

```

library(orthopolynom)

## Loading required package: polynom
DT$EnvN <- as.numeric(as.factor(DT$Env))
ansRR <- mmer(Yield~Env,
               random= ~ vs(leg(EnvN,1),Name),
               rcov= ~ units,
               data=DT)

## iteration      LogLik      wall      cpu(sec)      restrained
##     1      -40.232  21:52:13        0          0
##     2      -29.2803 21:52:13        0          0
##     3      -27.8646 21:52:13        0          0
##     4      -27.7107 21:52:13        0          0
##     5      -27.7036 21:52:13        0          0
##     6      -27.7032 21:52:13        0          0

summary(ansRR)

## =====
##           Multivariate Linear Mixed Model fit by REML
## ***** sommer 4.1 *****
## =====
##      logLik      AIC      BIC Method Converge
## Value -27.70318 61.40636 71.06743      NR      TRUE
## =====
## Variance-Covariance components:
##                               VarComp VarCompSE Zratio Constraint
## leg0:Name.Yield-Yield   10.392    3.1473  3.302  Positive
## leg1:Name.Yield-Yield   2.079    0.9792  2.123  Positive
## units.Yield-Yield       6.297    0.8442  7.459  Positive
## =====
## Fixed effects:
##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept) 16.541    0.6770  24.432
## 2 Yield EnvCA.2012   -5.832    0.6425 -9.078
## 3 Yield EnvCA.2013   -6.472    0.8239 -7.854
## =====

```

```

## Groups and observations:
##          Yield
## leg0:Name    41
## leg1:Name    41
## -----
## Use the '$' sign to access results and parameters

```

8) Other GxE covariance structures

A multi environment model is the one that is fitted when the breeding program can afford more than one location. This assumes that GxE doesn't exist and that the main genotype effect plus the fixed effect for environment is enough to predict the genotype effect in all locations of interest.

```

E <- CS(DT$Env)
rownames(E) <- colnames(E) <- unique(DT$Env)
EA <- kronecker(E,A, make.dimnames = TRUE)
ansCS <- mmr(Yield~Env,
              random= ~ vs(Name, Gu=A) + vs(Env:Name, Gu=EA),
              rcov= ~ units,
              data=DT)

## iteration      LogLik      wall      cpu(sec)      restrained
##   1      -31.1056  21:52:13        0            0
##   2      -23.39    21:52:13        0            0
##   3      -20.4917  21:52:13        0            0
##   4      -20.1504  21:52:13        0            0
##   5      -20.1454  21:52:13        0            0
##   6      -20.1454  21:52:13        0            0

summary(ansCS)

## =====
##           Multivariate Linear Mixed Model fit by REML
## ***** sommer 4.1 *****
## =====
##      logLik      AIC      BIC Method Converge
## Value -20.14538 46.29075 55.95182      NR      TRUE
## =====
## Variance-Covariance components:
##             VarComp VarCompSE Zratio Constraint
## u:Name.Yield-Yield     1.958     1.932  1.013  Positive
## u:Env:Name.Yield-Yield   6.897     1.994  3.460  Positive
## units.Yield-Yield       4.366     0.647  6.748  Positive
## =====
## Fixed effects:
##   Trait      Effect Estimate Std.Error t.value
## 1 Yield (Intercept)  16.496    0.6855  24.065
## 2 Yield EnvCA.2012   -5.777    0.7558  -7.643
## 3 Yield EnvCA.2013   -6.380    0.7960  -8.015
## =====
## Groups and observations:
##          Yield
## u:Name      41
## u:Env:Name  123
## =====

```

```
## Use the '$' sign to access results and parameters
```

Final remarks

Keep in mind that sommer uses direct inversion (DI) algorithm which can be very slow for large datasets. The package is focused in problems of the type $p > n$ (more random effect levels than observations) and models with dense covariance structures. For example, for experiment with dense covariance structures with low-replication (i.e. 2000 records from 1000 individuals replicated twice with a covariance structure of 1000x1000) sommer will be faster than MME-based software. Also for genomic problems with large number of random effect levels, i.e. 300 individuals (n) with 100,000 genetic markers (p). For highly replicated trials with small covariance structures or $n > p$ (i.e. 2000 records from 200 individuals replicated 10 times with covariance structure of 200x200) asreml or other MME-based algorithms will be much faster and we recommend you to opt for those software.

Literature

- Covarrubias-Pazaran G. 2016. Genome assisted prediction of quantitative traits using the R package sommer. PLoS ONE 11(6):1-15.
- Covarrubias-Pazaran G. 2018. Software update: Moving the R package sommer to multivariate mixed models for genome-assisted prediction. doi: <https://doi.org/10.1101/354639>
- Bernardo Rex. 2010. Breeding for quantitative traits in plants. Second edition. Stemma Press. 390 pp.
- Gilmour et al. 1995. Average Information REML: An efficient algorithm for variance parameter estimation in linear mixed models. Biometrics 51(4):1440-1450.
- Henderson C.R. 1975. Best Linear Unbiased Estimation and Prediction under a Selection Model. Biometrics vol. 31(2):423-447.
- Kang et al. 2008. Efficient control of population structure in model organism association mapping. Genetics 178:1709-1723.
- Lee, D.-J., Durban, M., and Eilers, P.H.C. (2013). Efficient two-dimensional smoothing with P-spline ANOVA mixed models and nested bases. Computational Statistics and Data Analysis, 61, 22 - 37.
- Lee et al. 2015. MTG2: An efficient algorithm for multivariate linear mixed model analysis based on genomic information. Cold Spring Harbor. doi: <http://dx.doi.org/10.1101/027201>.
- Maier et al. 2015. Joint analysis of psychiatric disorders increases accuracy of risk prediction for schizophrenia, bipolar disorder, and major depressive disorder. Am J Hum Genet; 96(2):283-294.
- Rodriguez-Alvarez, Maria Xose, et al. Correcting for spatial heterogeneity in plant breeding experiments with P-splines. Spatial Statistics 23 (2018): 52-71.
- Searle. 1993. Applying the EM algorithm to calculating ML and REML estimates of variance components. Paper invited for the 1993 American Statistical Association Meeting, San Francisco.
- Yu et al. 2006. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. Genetics 38:203-208.
- Abdollahi Arpanahi R, Morota G, Valente BD, Kranis A, Rosa GJM, Gianola D. 2015. Assessment of bagging GBLUP for whole genome prediction of broiler chicken traits. Journal of Animal Breeding and Genetics 132:218-228.
- Tunnicliffe W. 1989. On the use of marginal likelihood in time series model estimation. JRSS 51(1):15-27.